

Short Communication

Virtual Reality as a Treatment for Vestibular Dysfunction: A New Paradigm of Clinical Physiotherapy

Samanta Kariella Silva Figueiredo¹, Marco Orsini², Silmar Teixeira³, Bruna Velasques^{5,6}, Pedro Ribeiro^{5,6}, Nélcio Silva de Souza⁷, Marco Antônio A Leite⁸ and Victor Hugo do Vale Bastos^{4,5*}

¹Physiotherapist the Federal University of Piauí (UFPI), Brazil

²Master's Program in Rehabilitation Sciences (University Center of Augusto Motta – UNISUAM) and Collaborating Professor Masters/Doctoral Program in Neurology – Antônio Pedro University Hospital/Fluminense Federal University (HUAP/UFF), Brazil

³Brain Mapping and Plasticity Laboratory, Federal University of Piauí (LAMPLACE/ UFPI), Parnaíba, Brazil

⁴Brain Mapping and Functionality Laboratory (LAMCEF/ UFPI), Federal University of Piauí, Brazil

⁵Brain Mapping and Sensory Motor Integration, Institute of Psychiatry of Federal University of Rio de Janeiro (IPUB/UFRJ), Brazil

⁶School of Physical Education, Bioscience Department (EEFD/UFRJ), Brazil; Institute of Applied Neuroscience (INA), Rio de Janeiro, Brazil

⁷Physiotherapist, (University Center Serra dos Órgãos - UNIFESO); Master in Science of Rehabilitation (UNISUAM), Rio de Janeiro, RJ, Brazil

⁸Medical school (UFF); Neurology Division – Head of Movement Disorders Unit (HUAP/UFF), Brazil

*Corresponding author: Victor Hugo do Vale Bastos, Brain Mapping and Functionality Laboratory (LAMCEF/ UFPI), Federal University of Piauí, Brazil, Email: victorhugobastos@ufpi.edu.br

Received: June 15, 2015; Accepted: June 19, 2015;
Published: June 22, 2015

Short Communication

The vestibular system is a reference in balance control because it acts as a gravity sensor controlling the dynamic posture when the individual is walking, for example [1, 2]. Vestibular dysfunctions may be peripheral and/or central, occurring by primary or secondary causes, having as main manifestations, body sway under visual or somatosensory conflict conditions, reduced stability and functional capacity, gait deviations and falls [3]. Vestibular rehabilitation is based on mechanisms related to neuronal plasticity of the central nervous system, known as adaptation, habituation and substitution, where nerve cells establish new synaptic connections expanding its neural network [4]. A relatively new treatment based on games or virtual activities has been used in vestibular rehabilitation. Thus, the aim of this study is to present the effects of exposure therapy to virtual reality in vestibular rehabilitation.

Therapy in virtual reality suggests a connection between the individual and the machine, implying that it is in another reality in which the perception of the environment is modified by sensory

Abstract

The vestibular system is a reference in balance control because it acts as a gravity sensor controlling the dynamic posture. Vestibular dysfunctions may be peripheral and/or central, presents different manifestations (visual or somatosensory conflict), reduced stability and functional capacity, gait deviations and falls. The use of virtual reality has use in treatment of phobias, Parkinson's disease, after stroke, children with cerebral palsy as well as vestibular disorders. Vestibular rehabilitation is based on mechanisms related to neuronal plasticity of the central nervous system and virtual games activities have been used. The virtual reality therapy in the treatment of vestibular dysfunction presents few studies run on journals; however, the few studies have evidenced good results, which, nevertheless, need to be further explored, applying stricter protocols. The therapy has contributed to improvement of postural balance, motor coordination, flexibility, physical function, also increased stability limit, reduce dizziness levels and rates of falls, providing functional independence and quality of life. Considering that, this feature may be relevant in the treatment of vestibular dysfunctions it is essential to investigate how virtual reality is being aggregated in the vestibular rehabilitation process.

Keywords: Virtual reality; Vestibular dysfunction and physiotherapy

stimuli [5]. The use of virtual reality in healthcare was used since there was the first case of success in the treatment of phobias [6] and currently is being used in the rehabilitation of patients with Parkinson's disease, in the upper limb recovery of hemiparetic patients after stroke [7] and in the treatment of children with cerebral palsy with emphasis on motor control and balance [8].

In virtual reality exposure therapy it is possible to simulate computationally real spaces in a multisensory 3D context through resources that increase feelings such as special helmets, belts, headphones, platforms and equipment with LCD displays, that is, the therapy allows the user to interact with the virtual world in real time through multi sensations (hearing, sight, touch and smell) to feel involved and motivated to perform a given activity. The visible benefit of this therapeutic mode is the provision of a variety of stimuli if compared to traditional methods of vestibular rehabilitation presenting the patient sensory conflicts at different levels of difficulty and in a safe environment that will promote correction of balance and posture, improved mobility, the functionality of upper and lower limbs, besides promoting greater motivation for the patient in exercise performances [9-11].

The virtual reality therapy in the treatment of vestibular dysfunction presents few studies run on journals; however, the few studies have evidenced good results, which, nevertheless, need to be further explored, applying stricter protocols. The therapy has contributed to improvement of postural balance, motor coordination mobility, flexibility, physical function, also increased stability limit; reduce dizziness levels and rates of falls, providing functional independence and quality of life [5, 12].

Considering that, this feature may be relevant in the treatment of

vestibular dysfunctions it is essential to investigate how virtual reality is being aggregated in the vestibular rehabilitation process, what types of devices and games are being more used and how effective they are. So that the therapy for virtual reality exposure become customary in clinics, offices and even in the households.

References

1. Xerri C, Barthélémy J, Harlay F, Borel L, Lacour M. Neuronal coding of linear motion in the vestibular nuclei of the alert cat. I. Response characteristics to vertical otolith stimulation. *Exp Brain Res*. 1987; 65: 569-581.
2. Fitzpatrick R, McCloskey DI. Proprioceptive, visual and vestibular thresholds for the perception of sway during standing in humans. *J Physiol*. 1994; 478 : 173-186.
3. Quitschal RM, Fukunaga JY, Ganança MM, Caovilla HH. Evaluation of postural control in unilateral vestibular hypofunction. *Braz J Otorhinolaryngol*. 2014; 80: 339-345.
4. Brandt T, Strupp M, Dieterich M. Five keys for diagnosing most vertigo, dizziness, and imbalance syndromes: an expert opinion. *J Neurol*. 2014; 261: 229-231.
5. Cho GH, Hwangbo G, Shin HS. The Effects of Virtual Reality-based Balance Training on Balance of the Elderly. *J Phys Ther Sci*. 2014; 26: 615-617.
6. Castro W, Sánchez M, Sedeño T. Los nuevos desarrollos tecnológicos aplicados al tratamiento psicológico. *Acta Colombiana de Psicología*. 2014; 17: 91-101.
7. Soares A, Woellner S, Andrade C, Mesadri T, Bruckheimer A, Hounsell M. The use of Virtual Reality for upper limb rehabilitation of hemiparetic Stroke patients. *Fisiot em Movim*. 2014; 27: 309-317.
8. Barela JA, Focks GM, Hilgeholt T, Barela AM, Carvalho Rde P, Savelsbergh GJ. Perception-action and adaptation in postural control of children and adolescents with cerebral palsy. *Res Dev Disabil*. 2011; 32: 2075-2083.
9. Garcia AP, Ganança MM, Cusin FS, Tomaz A, Ganança FF, Caovilla HH. Vestibular rehabilitation with virtual reality in Ménière's disease. *Braz J Otorhinolaryngol*. 2013; 79: 366-374.
10. Meldrum D, Herdman S, Moloney R, Murray D, Duffy D, Malone K, et al. Effectiveness of conventional versus virtual reality based vestibular rehabilitation in the treatment of dizziness, gait and balance impairment in adults with unilateral peripheral vestibular loss: a randomised controlled trial. *BMC Ear, Nose and Throat Disorders*. 2012; 12: 3.
11. Whitney SL, Sparto PJ, Brown KE, Furman JM, Jacobson JL, Redfern MS. The potential use of virtual reality in vestibular rehabilitation: preliminary findings with the BNAVE. *J Neurol Phys Ther*. 2002; 26: 72-78.
12. Macedo C, Gazzola JM, Ricci NA, Doná F, Ganança FF. Influence of sensory information on static balance in older patients with vestibular disorder. *Braz J Otorhinolaryngol*. 2015; 81: 50-57.